Semiclassical Approximations And Predictability In Ocean Acoustics

Michael G. Brown
Division of Applied Marine Physics
Rosenstiel School of Marine and Atmospheric Science
University of Miami
4600 Rickenbacker Causeway
Miami, FL 33149-1098

phone: (305) 361-4640 fax: (305) 361-4701 email: mbrown@rsmas.miami.edu

Award #: N000149710046

http://www.rsmas.miami.edu/divs/amp.html

LONG-TERM GOAL

Our long-term goal is to improve our understanding of the physics of the forward scattering of underwater sound. Of particular interest is understanding limitations on the predictability of underwater sound fields, especially using semiclassical (ray-based) methods. Improving our understanding of the mechanism(s) that contribute to the loss of predictability will lead to improved predictive models.

OBJECTIVES

The scientific objective of this work is to understand limitations on the predictability of acoustic wavefields in realistic (range-dependent) ocean environments. Both full wave and semiclassical predictive models are of interest. Both cw and broadband wavefields are of interest. We seek to both develop improved predictive acoustic models and to understand their limitations.

APPROACH

Using semiclassical methods, ideas relating to ray chaos provide a framework for studying predictability. Wavefield behavior in the ray limit is explored numerically and, whenever possible, analytically. The extent to which limitations on predictability carry over to finite frequency wavefields are then explored numerically, generally with parabolic equation based models. Both deterministic and stochastic models are of interest. Comparisons with data are made whenever possible.

Several tools have recently been developed and/or extended for use in our work: Tappert's coinsensitive PE model; the MaChl algorithm (Brown, 1994) for finite frequency ray-based wavefield predictions; a new stochastic ray model (Brown and Viechnicki, 1998) to model scattering by internal waves; and a new efficient algorithm to generate realistic internal wave induced sound speed perturbation fields (Colosi and Brown, 1998).

This work is being done in loose collaboration with the following individuals: F. Tappert (U. Miami; parabolic wave equations, waves in random media, ray chaos); M. Wolfson (Washington State U.; numerical modelling, waves in random media, ray chaos); G. Zaslavsky (Courant Institute; chaos in dynamical systems, stochastic methods, quantum chaos); S. Tomsovic (Washington State U.; quantum chaos and semiclassical breakdown); and J. Colosi (WHOI; internal waves, long-range propagation).

maintaining the data needed, and coincluding suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu ald be aware that notwithstanding an DMB control number.	ion of information. Send comments arters Services, Directorate for Info	s regarding this burden estimate ormation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 1998 2. REPOR		2. REPORT TYPE	3. DATES COVERED 00-00-1998 to 00-00-1998		
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
Semiclassical Approximations and Predictability in Ocean Acoustics				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Miami,Rosenstiel School of Marine and Atmospheric Science,4600 Rickenbacker Causeway,Miami,FL,33149				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAIL Approved for publ	ABILITY STATEMENT ic release; distributi	ion unlimited			
13. SUPPLEMENTARY NO See also ADM0022					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	3	

Report Documentation Page

Form Approved OMB No. 0704-0188

WORK COMPLETED

The principal accomplishment of the recent years are 1) the development and numerical implementation of stochastic ray theory to describe and model the scattering of sound by internal waves in deep ocean environments; 2) the development (with J. Colosi at WHOI) of a new efficient, accurate, and robust algorithm to compute internal-wave-induced sound speed perturbation fields; 3) incorporation of this internal wave model into both ray and full wave propagation models; 4) demonstration that for a large class of problems involving boundary interactions, ray phase errors are inversely proportional to frequency and increase linearly with range; and 5) demonstration (Brown, 1998) that, in the ray limit, islands of stability in phase space exist when the environment is almost stratified and has quasiperiodic range dependence.

RESULTS

Stochastic ray theory and the new technique to compute internal-wave-induced sound speed perturbation fields are important new tools which can be used to address fundamental questions relating to the predictability of long-range sound propagation in deep ocean environments. The work relating to the structure of phase space provides critically important insight into - and a mathematical framework for the study of - wave propagation in environments with nontrivial range-dependence. Our predictability studies show that, even in the presence of chaotic ray motion, semiclassical wavefield representations accurately predict many important features of underwater acoustic wavefields.

IMPACT/APPLICATION

Our work gives insight into the limitations on the predictability of underwater sound fields. This is an important basic science issue which impacts all systems applications which require accurate predictions of underwater sound fields.

TRANSITIONS

The PI collaborates informally with the investigators listed above, ATOC investigators, and others. This includes the sharing of both ideas and software. It is not known whether any software produced by the PI has been used to address any applied Navy problems.

RELATED PROJECTS

This work is closely related to the ATOC project and the ONR-funded work being performed by P. Worcester (SIO), J. Colosi (WHOI), M. Wolfson (WSU), J. Spiesberger (PSU), S. Tomsovic (WSU), G. Zaslavsky (CIMS) and F. Tappert (U. Miami). All of these projects are concerned with aspects of long-range propagation.

REFERENCES

Brown, M.G., 1994, A Maslov-Chapman wavefield representation for wide-angle one-way propagation, Geophys. J. Intl. 116, 513-5263.

Brown, M.G., and J. Viechnicki, 1998, Stochastic ray theory for long-range sound propagation in deep ocean environments, J. Acoust. Soc. Am. 104, 2090-2104.

Colosi, J.A., and M.G. Brown, 1998, Efficient numerical simulation of stochastic internal wave induced sound speed perturbation fields, J. Acoust. Soc. Am. 103, 2232-2235.

Browm, M.G., 1998, Phase space structure and fractal trajectories in 1 1/2 degree of freedom Hamiltonian systems whose time dependence is quasiperiodic, Nonlinear Processes in Geophysics, in press.

PUBLICATIONS

Brown, M.G., and J. Viechnicki, 1998, Stochastic ray theory for long-range sound propagation in deep ocean environments, J. Acoust. Soc. Am. 104, 2090-2104.

Colosi, J.A., and M.G. Brown, 1998, Efficient numerical simulation of stochastic internal wave induced sound speed perturbation fields, J. Acoust. Soc. Am. 103, 2232-2235.

Browm, M.G., 1998, Phase space structure and fractal trajectories in 1 1/2 degree of freedom Hamiltonian systems whose time dependence is quasiperiodic, Nonlinear Processes in Geophysics, in press.